

Exchange Relations for the q -vertex operators of $U_q(\widehat{sl_2})$

HIDETOSHI AWATA ¹

*Dept. of Physics Hokkaido University,
Sapporo 060, Japan*

Abstract

We consider the q -deformed Knizhnik-Zamolodchikov equation for the two point function of q -deformed vertex operators of $U_q(\widehat{sl_2})$. We give explicitly the fundamental solutions, the connection matrices and the exchange relations for the q -vertex operators of spin $1/2$ and $j \in \frac{1}{2}\mathbf{Z}_{\geq 0}$. Consequently, we confirm that the connection matrices are equivalent to the elliptic Boltzman weights of IRF type obtained by the fusion procedure from ABF models.

hep-th/9211087

¹ E-mail address : hawa@theory.kek.jp

1. Introduction

Recently, the q -deformed Knizhnik-Zamolodchikov equation (q -KZ eq.), more generally holonomic q -difference equation, has been analyzed [1,2,3,4,5,6] and the remarkable relation between the connection matrices of its solutions and the elliptic Boltzman weights was investigated [4,7].

To study the q -KZ equation², an important and useful concept is the q -vertex operators, which are q -analogue of the vertex operators in Ref. [10]. First, the solutions of the q -KZ equation can be constructed as the correlation functions of the q -vertex operators [3,4]. Moreover, we can understand naturally the IRF type Yang-Baxter relation satisfied by the connection matrices of q -KZ solutions as the exchange relation for the q -vertex operators [4,11]. So far, explicit calculation of the exchange relation for the q -vertex operators has been done in Ref. [4] for the first nontrivial examples and they conjectured that the connection matrices are generally equivalent to the Boltzman weights in Ref. [12,13]. The exchange relations of the q -vertex operators of spin $1/2$ or $k/2$ (of level k) were calculated and applied to the vertex models in Ref. [8,9].

The aim of this paper is to generalize these results to arbitrary spins. We consider the q -KZ equation for the two point function of the q -vertex operators of spin $1/2$ and $j \in \frac{1}{2}\mathbf{Z}_{\geq 0}$ and we give the fundamental solutions and the connection matrices explicitly.

2. Quantum affine algebra $U_q(\widehat{sl_2})$

§ 2.1. First we fix some notation. The algebra $U_q(\widehat{sl_2})$ is generated by e_i , f_i , invertible k_i ($i = 0, 1$) and d with relations

$$\begin{aligned} k_i e_j k_i^{-1} &= q^{a_{ij}} e_j, & \sum_{n=0}^3 (-1)^n \begin{bmatrix} 3 \\ n \end{bmatrix} e_i^{3-n} e_j e_i^n &= 0, & [d, e_i] &= \delta_{i,0} e_i, \\ k_i f_j k_i^{-1} &= q^{-a_{ij}} f_j, & \sum_{n=0}^3 (-1)^n \begin{bmatrix} 3 \\ n \end{bmatrix} f_i^{3-n} f_j f_i^n &= 0, & [d, k_i] &= 0, \\ e_i f_j - f_j e_i &= \delta_{ij} \frac{k_i - k_i^{-1}}{q - q^{-1}}, & & & [d, f_i] &= -\delta_{i,0} f_i, \end{aligned} \quad (2.1)$$

² The q -KZ equation has been applied to the Thirring model in Ref. [3] and to the XXZ model or the vertex model in Ref. [8,9].

and $k_0 k_1 = q^k$ with a level $k \in \mathbf{C}$, where $q \in \mathbf{C}$, $a_{11} = a_{00} = -a_{10} = -a_{01} = 2$ and

$$\begin{bmatrix} n \\ m \end{bmatrix} = \frac{[n]!}{[n-m]![m]!}, \quad [n] = \frac{q^n - q^{-n}}{q - q^{-1}}. \quad (2.2)$$

The algebra $U_q(\widehat{sl_2})$ is a Hopf algebra with the comultiplication Δ , the antipode S and the co-unit ϵ

$$\begin{aligned} \Delta(e_i) &= e_i \otimes k_i + 1 \otimes e_i, & S(e_i) &= -e_i k_i^{-1}, & \epsilon(e_i) &= 0, \\ \Delta(k_i) &= k_i \otimes k_i, & S(k_i) &= k_i^{-1}, & \epsilon(k_i) &= 1, \\ \Delta(f_i) &= f_i \otimes 1 + k_i^{-1} \otimes f_i, & S(f_i) &= -k_i f_i, & \epsilon(f_i) &= 0, \\ \Delta(d) &= d \otimes 1 + 1 \otimes d, & S(d) &= -d, & \epsilon(d) &= 0. \end{aligned} \quad (2.3)$$

§ 2.2. Let V_j be the Verma module over $U_q(\widehat{sl_2})$, generated by the highest weight vector $|j\rangle$, such that $e_i|j\rangle = 0$, $k_1|j\rangle = q^{2j}|j\rangle$ and $d|j\rangle = -h_j|j\rangle$ with $h_j = j(j+1)/\kappa$, $\kappa = k+2$. The dual module V_j^* is generated by $\langle j|$ which satisfies $\langle j|e_i = 0$, $\langle j|k_1 = q^{2j}\langle j|$ and $\langle j|d = -h_j\langle j|$. The bilinear form $V_j^* \otimes V_j \rightarrow \mathbf{C}$ is uniquely defined by $\langle j|j\rangle = 1$ and $(\langle u|a)|v\rangle = \langle u|(a|v\rangle)$ for any $\langle u| \in V_j^*$, $|v\rangle \in V_j$ and $a \in U_q(\widehat{sl_2})$.

A null vector $|\chi\rangle \in V_j$ (of grade N and charge Q) is defined by $e_i|\chi\rangle = 0$, $k_1|\chi\rangle = q^{2(j+Q)}|\chi\rangle$ and $d|\chi\rangle = -(h_j + N)|\chi\rangle$. A null vector $\langle\chi| \in V_j^*$ is defined in a similar manner.

For $2j \in \mathbf{Z}_{\geq 0}$, we have a $2j+1$ dimensional centerless irreducible representation of $U_q(\widehat{sl_2})$, $V_j(z) = \oplus_{m=0}^{2j} \mathbf{C}(q, z)|j, m\rangle$, which is defined by

$$\begin{aligned} e_1|j, m\rangle &= [2j - m + 1]|j, m-1\rangle, & e_0|j, m\rangle &= z[m+1]|j, m+1\rangle, \\ k_1|j, m\rangle &= q^{2(j-m)}|j, m\rangle, & k_0|j, m\rangle &= q^{-2(j-m)}|j, m\rangle, \\ f_1|j, m\rangle &= [m+1]|j, m+1\rangle, & f_0|j, m\rangle &= z^{-1}[2j - m + 1]|j, m-1\rangle, \end{aligned} \quad (2.4)$$

and d acts as $d = -h_j + z \frac{d}{dz}$.

3. The q -vertex operators and their two point functions

§ 3.1. The q -vertex operator, $\Phi_{j_2}(z) : V_{j_1} \rightarrow V_{j_3}$, of spin $j_2 \in \mathbf{C}$ is defined by the transformation property under the adjoint action [4,3,14], or equivalently as an intertwiner (see Appendix A).

Definition. For $2j_2 \in \mathbf{Z}_{\geq 0}$, the q -vertex operator ³, $\Phi_{j_2, m_2}^{\alpha_2}(z) : V_{j_1} \rightarrow V_{j_3}$ with $j_3 = j_1 + j_2 - \alpha_2$ and $0 \leq m_2 \leq 2j_2$, is defined explicitly as follows ⁴

$$\begin{aligned}
e_1 \Phi_{j, m}(z) &= [2j - m + 1] \Phi_{j, m-1}(z) k_1 + \Phi_{j, m}(z) e_1, \\
k_1 \Phi_{j, m}(z) &= q^{2(j-m)} \Phi_{j, m}(z) k_1, \\
f_1 \Phi_{j, m}(z) &= [m + 1] \Phi_{j, m+1}(z) + q^{-2(j-m)} \Phi_{j, m}(z) f_1, \\
e_0 \Phi_{j, m}(z) &= z[m + 1] \Phi_{j, m+1}(z) k_0 + \Phi_{j, m}(z) e_0, \\
k_0 \Phi_{j, m}(z) &= q^{-2(j-m)} \Phi_{j, m}(z) k_0, \\
f_0 \Phi_{j, m}(z) &= z^{-1}[2j - m + 1] \Phi_{j, m-1}(z) + q^{2(j-m)} \Phi_{j, m}(z) f_0,
\end{aligned} \tag{3.1}$$

and ⁵

$$d \Phi_{j, m}(z) = -z \frac{d}{dz} \Phi_{j, m}(z) + \Phi_{j, m}(z) d. \tag{3.2}$$

The existence conditions for the q -vertex operator $\Phi_{j, m}(z)$ with a general spin $j \in \mathbf{C}$ were analyzed in Ref. [15], and we will review them in Appendix A. In the case that q is not a root of unity, the existence conditions are essentially the same as those of $q = 1$ [16]. For integrable representations of general quantum affine algebras, the complete results on existence and uniqueness were given in Ref. [11].

From the k_1 and d commutation relations, the ground state matrix element of the q -vertex operator can be determined up to normalization, and we normalize it as

$$\langle j_3 | \Phi_{j_2, m_2}^{\alpha_2}(z) | j_1 \rangle = \delta_{j_2 - m_2}^{j_3 - j_1} \delta_{j_2 - \alpha_2}^{j_3 - j_1} z^{h_3 - h_1}, \tag{3.3}$$

³ Our q -vertex operators correspond to the type-II vertex operators in Ref. [11, 8, 9,]. The relation between our q -vertex operator $\Phi_{j_2, m_2}^{j_1 + j_2 - j_3}(z) : V_{j_1} \rightarrow V_{j_3}$ and their one $\Phi_{j_2, j_1}^{j_3}(z) : V_{j_2}(z) \otimes V_{j_1} \rightarrow V_{j_3}$ is

$$\Phi_{j_2, m_2}^{j_1 + j_2 - j_3}(z) = \Phi_{j_2, j_1}^{j_3}(z) \left(|j_2, m_2\rangle, * \right).$$

The $\Phi_{j_2, j_1}^{j_3}(z)$ has the intertwining property, $a \Phi_{j_2, j_1}^{j_3}(z) = \Phi_{j_2, j_1}^{j_3}(z) \Delta(a)$, for all $a \in U_q(\widehat{sl_2})$. Note that the coproduct is also slightly different.

⁴ The upper index α , which specify the modules on which the q -vertex operator acts, will be sometimes suppressed.

⁵ The sign difference of d in §2.2 and §3.1 comes from the difference in meaning of z . In §2.2, z is a generator such that $[d, z] = 1$, on the other hand z is just a variable i.e. $[d, z] = 0$, in §3.1.

where $h_n = h_{j_n}$. The other matrix elements for the descendant fields can be uniquely determined.

We can in principle derive the arbitrary N point functions by using this one point function and the q -operator product expansion (q -OPE). In Appendix B, we will present the q -OPE of a spin $1/2$ q -vertex operator and the two point functions $\langle j_4 | \Phi_{\frac{1}{2}, m_3}(z_3) \Phi_{j_2, m_2}(z_2) | j_1 \rangle$ and $\langle j_4 | \Phi_{j_2, m_2}(z_2) \Phi_{\frac{1}{2}, m_3}(z_3) | j_1 \rangle$.

§ **3.2.** As we will discuss in Appendix B, the above two point functions have a complicated form, for example each coefficient of z^n can not be factorised. But they can be simplified by dividing by a function

$$g(x) = (1 - q^{k+2} \frac{[2j]}{[2][k+2]} x + O(x^2)), \quad x = z_2/z_3. \quad (3.4)$$

Here we list the leading terms

$$\begin{aligned} \langle j_4 | \Phi_{\frac{1}{2}, 0}^0(z_3) \Phi_{j_2, M}^M(z_2) | j_1 \rangle &= z_3^{h_4 - h_4^-} z_2^{h_4^- - h_1} \\ &\times \{1 + p^{-m_1 - m_2} x \frac{[2m + m_1 + m_2 + 1]_p [2m_1]_p}{[2m + m_1 - m_2 + 1]_p} + O(x^2)\} g(x), \\ \langle j_4 | \Phi_{\frac{1}{2}, 1}^0(z_3) \Phi_{j_2, M-1}^M(z_2) | j_1 \rangle &= -z_3^{h_4 - h_4^-} z_2^{h_4^- - h_1} x p^{-m_2} \frac{[2m_1]_p}{[2m + m_1 - m_2 + 1]_p} \\ &\times \{1 + p^{-m_1 - m_2} x \frac{[2m + m_1 + m_2 + 1]_p [2m_1 + 1]_p}{[2m + m_1 - m_2 + 2]_p} + O(x^2)\} g(x), \end{aligned} \quad (3.5)$$

$$\begin{aligned} \langle j_4 | \Phi_{\frac{1}{2}, 0}^1(z_3) \Phi_{j_2, M}^{M-1}(z_2) | j_1 \rangle &= -z_3^{h_4 - h_4^+} z_2^{h_4^+ - h_1} p^{-m_1} \frac{[2m_2]_p}{[-2m - m_1 + m_2]_p} \\ &\times \{1 + p^{-m_1 - m_2} x \frac{[-2m + m_1 + m_2]_p [2m_2 + 1]_p}{[-2m - m_1 + m_2 + 1]_p} + O(x^2)\} g(x), \end{aligned} \quad (3.6)$$

$$\begin{aligned} \langle j_4 | \Phi_{\frac{1}{2}, 1}^1(z_3) \Phi_{j_2, M-1}^{M-1}(z_2) | j_1 \rangle &= z_3^{h_4 - h_4^+} z_2^{h_4^+ - h_1} \\ &\times \{1 + p^{-m_1 - m_2} x \frac{[-2m + m_1 + m_2]_p [2m_2]_p}{[-2m - m_1 + m_2]_p} + O(x^2)\} g(x), \end{aligned}$$

and

$$\begin{aligned}
\langle j_4 | \Phi_{j_2, M}^M(z_2) \Phi_{\frac{1}{2}, 0}^0(z_3) | j_1 \rangle &= z_2^{h_4 - h_1^+} z_3^{h_1^+ - h_1} \\
&\times \{1 + p^{-m_1 - m_2} x^{-1} \frac{[-2m + m_1 + m_2]_p [2m_1]_p}{[-2m + m_1 - m_2]_p} + O(x^{-2})\} g(x^{-1}), \\
\langle j_4 | \Phi_{j_2, M-1}^M(z_2) \Phi_{\frac{1}{2}, 1}^0(z_3) | j_1 \rangle &= -z_2^{h_4 - h_1^+} z_3^{h_1^+ - h_1} p^{-m_2} \frac{[2m_1]_p}{[-2m + m_1 - m_2]_p} \\
&\times \{1 + p^{-m_1 - m_2} x^{-1} \frac{[-2m + m_1 + m_2]_p [2m_1 + 1]_p}{[-2m + m_1 - m_2 + 1]_p} + O(x^{-2})\} g(x^{-1}), \\
\langle j_4 | \Phi_{j_2, M}^{M-1}(z_2) \Phi_{\frac{1}{2}, 0}^1(z_3) | j_1 \rangle &= -z_2^{h_4 - h_1^-} z_3^{h_1^- - h_1} x^{-1} p^{-m_1} \frac{[2m_2]_p}{[2m - m_1 + m_2 + 1]_p} \\
&\times \{1 + p^{-m_1 - m_2} x^{-1} \frac{[2m + m_1 + m_2 + 1]_p [2m_2 + 1]_p}{[2m - m_1 + m_2 + 2]_p} + O(x^{-2})\} g(x^{-1}), \\
\langle j_4 | \Phi_{j_2, M-1}^{M-1}(z_2) \Phi_{\frac{1}{2}, 1}^1(z_3) | j_1 \rangle &= z_2^{h_4 - h_1^-} z_3^{h_1^- - h_1} \\
&\times \{1 + p^{-m_1 - m_2} x^{-1} \frac{[2m + m_1 + m_2 + 1]_p [2m_2]_p}{[2m - m_1 + m_2 + 1]_p} + O(x^{-2})\} g(x^{-1}),
\end{aligned} \tag{3.7}$$

$$\begin{aligned}
&\times \{1 + p^{-m_1 - m_2} x^{-1} \frac{[2m + m_1 + m_2 + 1]_p [2m_2]_p}{[2m - m_1 + m_2 + 1]_p} + O(x^{-2})\} g(x^{-1}), \\
&\times \{1 + p^{-m_1 - m_2} x^{-1} \frac{[2m + m_1 + m_2 + 1]_p [2m_2]_p}{[2m - m_1 + m_2 + 1]_p} + O(x^{-2})\} g(x^{-1}),
\end{aligned} \tag{3.8}$$

here and below we use the following notations, $[n]_p = (p^n - 1)/(p - 1)$, $p = q^{-2\kappa}$, $m = -(j_1 + j_4 + 1)/2\kappa$, $m_1 = (j_1 + j_2 - j_4 + \frac{1}{2})/2\kappa = M/2\kappa$, $m_2 = (-j_1 + j_2 + j_4 + \frac{1}{2})/2\kappa = \overline{M}/2\kappa$ and $h_n^\pm = h_{j_n \pm \frac{1}{2}}$.

More complete expressions for the two point functions will be given in §5.1 by solving the q -KZ equation.

4. The solutions and their connection formula for the q -KZ equation

§ 4.1. Arbitrary N point functions of q -vertex operators satisfy the q -KZ equation [4]. For the two point function of the q -vertex operators of spin $1/2$ and j , $\langle j_4 | \Phi_{\frac{1}{2}, M_3}^1(z_3) \Phi_{j_2, M_2}^1(z_2) | j_1 \rangle$, the q -KZ equation is written as a 2×2 block diagonal R matrix $R(z_2/z_3) : V_{j_2}(z_2) \otimes V_{\frac{1}{2}}(z_3) \rightarrow V_{j_2}(z_2) \otimes V_{\frac{1}{2}}(z_3)$, defined by $R(z_2/z_3) \Delta'_{z_2, z_3} = \Delta_{z_2, z_3} R(z_2/z_3)$ (Appendix C).

Let $M = M_2 + M_3$, $M + \overline{M} = 2j_2 + 1$ and $x = z_2/z_3$.

Up to normalization, this q -KZ equation is

$$\begin{aligned} \begin{pmatrix} p^{-m} \tilde{\Psi}_0(px) \\ p^m \tilde{\Psi}_1(px) \end{pmatrix} &= \tilde{R}(x) \begin{pmatrix} \tilde{\Psi}_0(x) \\ \tilde{\Psi}_1(x) \end{pmatrix}, \\ \tilde{R}(x) &= \begin{pmatrix} \tilde{R}_0^0(x) & \tilde{R}_0^1(x) \\ \tilde{R}_1^0(x) & \tilde{R}_1^1(x) \end{pmatrix} = \frac{1}{1 - xp^{m_1+m_2}} \begin{pmatrix} p^{m_1} - xp^{m_2} & p^{m_2}(p^{-m_2} - p^{m_2}) \\ xp^{m_1}(p^{-m_1} - p^{m_1}) & p^{m_2} - xp^{m_1} \end{pmatrix}, \end{aligned} \quad (4.1)$$

where p , m and m_i are the same as in §3.2.

The equation (4.1) can be solved easily, and the solutions are given by the q -hypergeometric function

$$F_p(a, b, c; x) = \sum_{n=0}^{\infty} \frac{(a)_n (b)_n}{(c)_n (1)_n} x^n, \quad (4.2)$$

where $(a)_n = [a]_p [a+1]_p \cdots [a+n-1]_p$.

Two fundamental solutions $\tilde{\Psi}_{(+)}^0(x)$ and $\tilde{\Psi}_{(+)}^1(x)$ in the region $x \ll 1$ are

$$\begin{aligned} \tilde{\Psi}_{(+)}^0(x) &= -F(-2m + m_1 + m_2, 2m_2 + 1, -2m - m_1 + m_2 + 1; p^{-m_1-m_2}x) \\ &\quad \times p^{-m_1} \frac{[2m_2]_p}{[-2m - m_1 + m_2]_p} x^{m_2-m}, \\ \tilde{\Psi}_{(+)}^1(x) &= F(-2m + m_1 + m_2, 2m_2, -2m - m_1 + m_2; p^{-m_1-m_2}x) x^{m_2-m}, \\ \tilde{\Psi}_{(+)}^0(x) &= F(2m + m_1 + m_2 + 1, 2m_1, 2m + m_1 - m_2 + 1; p^{-m_1-m_2}x) x^{m+m_1}, \\ \tilde{\Psi}_{(+)}^1(x) &= -F(2m + m_1 + m_2 + 1, 2m_1 + 1, 2m + m_1 - m_2 + 2; p^{-m_1-m_2}x) \\ &\quad \times p^{-m_2} \frac{[2m_1]_p}{[2m + m_1 - m_2 + 1]_p} x^{m+m_1+1}. \end{aligned} \quad (4.3)$$

And the other two fundamental solutions $\tilde{\Psi}_{(-)}^0(x)$ and $\tilde{\Psi}_{(-)}^1(x)$ in the region $x \gg 1$ are

$$\begin{aligned} \tilde{\Psi}_{(-)}^0(x) &= -F(2m + m_1 + m_2 + 1, 2m_2 + 1, 2m - m_1 + m_2 + 2; p^{-m_1-m_2+1}x^{-1}) \\ &\quad \times p^{2m-m_1+1} \frac{[2m_2]_p}{[2m - m_1 + m_2 + 1]_p} x^{-m-m_2-1}, \\ \tilde{\Psi}_{(-)}^1(x) &= F(2m + m_1 + m_2 + 1, 2m_2, 2m - m_1 + m_2 + 1; p^{-m_1-m_2+1}x^{-1}) x^{-m-m_2}, \\ \tilde{\Psi}_{(-)}^0(x) &= F(-2m + m_1 + m_2, 2m_1, -2m + m_1 - m_2; p^{-m_1-m_2+1}x^{-1}) x^{m-m_1}, \\ \tilde{\Psi}_{(-)}^1(x) &= -F(-2m + m_1 + m_2, 2m_1 + 1, -2m + m_1 - m_2 + 1; p^{-m_1-m_2+1}x^{-1}) \\ &\quad \times p^{-2m-m_2} \frac{[2m_1]_p}{[-2m + m_1 - m_2]_p} x^{m-m_1}. \end{aligned} \quad (4.4)$$

§ 4.2. The complete q -KZ equation with the correct normalization is

$$\begin{pmatrix} p^{-m}\Psi_0(px) \\ p^m\Psi_1(px) \end{pmatrix} = R(x) \begin{pmatrix} \Psi_0(x) \\ \Psi_1(x) \end{pmatrix}, \quad (4.5)$$

where $R(x)$ is the image of the universal R -matrix which satisfies the crossing relation (see Appendix C). This $R(x)$ is given by

$$R(x) = f(x)\tilde{R}(x), \quad f(x) = \prod_{n \geq 0} \frac{(1 - xq^{2j_2+3+4n})(1 - xq^{-2j_2+1+4n})}{(1 - xq^{2j_2+1+4n})(1 - xq^{-2j_2+3+4n})}, \quad (4.6)$$

in the region $x \ll 1$.

The relation of the $\Psi(x)$'s and the previous $\tilde{\Psi}(x)$'s is

$$\Psi_i(x) = g(x)\tilde{\Psi}_i(x), \quad g(xp) = f(x)g(x). \quad (4.7)$$

The factor $g(x)$ in the regions $x \ll 1$ and $x \gg 1$ are

$$\begin{aligned} g_+(x) &= \prod_{n,m \geq 0} \frac{(1 - xq^{2j_2+1+4n}p^m)(1 - xq^{-2j_2+3+4n}p^m)}{(1 - xq^{2j_2+3+4n}p^m)(1 - xq^{-2j_2+1+4n}p^m)}, \\ g_-(x) &= \prod_{n,m \geq 0} \frac{(1 - x^{-1}q^{-2j_2-3-4n}p^{m+1})(1 - x^{-1}q^{2j_2-1-4n}p^{m+1})}{(1 - x^{-1}q^{-2j_2-1-4n}p^{m+1})(1 - x^{-1}q^{2j_2-3-4n}p^{m+1})}, \end{aligned} \quad (4.8)$$

respectively.

§ 4.3. The connection formula for the q -hypergeometric function $F_p(a, b, c; x)$ is ⁶

$$\begin{aligned} F_p(a, b, c; x) &= \frac{\Gamma_p(c)\Gamma_p(b-a)\Theta(p^a x, p)}{\Gamma_p(b)\Gamma_p(c-a)\Theta(x, p)} F_p(a, a-c+1, a-b+1, p^{c+1-a-b}x^{-1}) \\ &+ \frac{\Gamma_p(c)\Gamma_p(a-b)\Theta(p^b x, p)}{\Gamma_p(a)\Gamma_p(c-b)\Theta(x, p)} F_p(b, b-c+1, a-b+1, p^{c+1-a-b}x^{-1}), \end{aligned} \quad (4.9)$$

where

$$\begin{aligned} \Gamma_p(a) &= (1-p)^{1-a} \prod_{n \geq 0} \frac{(1-p^{n+1})}{(1-p^{n+a})}, \\ \Theta(x, p) &= \prod_{n \geq 0} (1-p^{n+1})(1-xp^n)(1-x^{-1}p^{n+1}). \end{aligned} \quad (4.10)$$

⁶ For the proof of this formula, see Ref. [2] for example.

Using this formula, we have

Proposition I. *The connection formulas for the solutions of the q -KZ equation (4.1) and for the normalization factor $g(x)$ are*

$$\tilde{\Psi}_{(+)}^\alpha(x) = \sum_{\beta} \tilde{\Psi}_{(-)}^\beta(x) C_\beta^\alpha(x), \quad (4.11)$$

$$\begin{aligned} C_0^0(x) &= \frac{\Gamma_p(2m+m_1-m_2+1)\Gamma_p(2m-m_1+m_2+1)}{\Gamma_p(2m-m_1-m_2+1)\Gamma_p(2m+m_1+m_2+1)} \frac{\Theta(p^{m_1-m_2}x)}{\Theta(p^{-m_1-m_2}x)} x^{2m_1}, \\ C_1^0(x) &= -\frac{\Gamma_p(-2m+m_1-m_2)\Gamma_p(2m+m_1-m_2+1)}{\Gamma_p(-2m_2+1)\Gamma_p(2m_1)} \frac{\Theta(p^{2m+1}x)}{\Theta(p^{-m_1-m_2}x)} p^{-m_2} x^{2m+m_1+m_2+1}, \\ C_0^1(x) &= -\frac{\Gamma_p(-2m-m_1+m_2)\Gamma_p(2m-m_1+m_2+1)}{\Gamma_p(-2m_1+1)\Gamma_p(2m_2)} \frac{\Theta(p^{-2m}x)}{\Theta(p^{-m_1-m_2}x)} p^{-m_1} x^{-2m+m_1+m_2}, \\ C_1^1(x) &= \frac{\Gamma_p(-2m+m_1-m_2)\Gamma_p(-2m-m_1+m_2)}{\Gamma_p(-2m-m_1-m_2)\Gamma_p(-2m+m_1+m_2)} \frac{\Theta(p^{m_2-m_1}x)}{\Theta(p^{-m_1-m_2}x)} x^{2m_2}, \end{aligned} \quad (4.12)$$

and

$$g_+(x) = g_-(x) C_g(x), \quad C_g(x) = \prod_{n \geq 0} \frac{\Theta(xq^{2j_2+1+4n})\Theta(xq^{-2j_2+3+4n})}{\Theta(xq^{2j_2+3+4n})\Theta(xq^{-2j_2+1+4n})}. \quad (4.13)$$

Note that $C_\beta^\alpha(x)$ and $C_g(x)$ are the pseudo-constant, e.g. $C_g(px) = C_g(x)$.

5. Exchange relation for the q -vertex operators

§ 5.1. Comparing the two point function in §3.2 and $\tilde{\Phi}(x)$ in §4.1, we have

Proposition II. *The relations between the solutions of the reduced q -KZ equation in §4.1 with the two point functions in §3.2 are*

$$\begin{aligned} & \begin{pmatrix} \langle j_4 | \Phi_{\frac{1}{2},0}^0(z_3) \Phi_{j_2,M}^M(z_2) | j_1 \rangle & \langle j_4 | \Phi_{\frac{1}{2},0}^1(z_3) \Phi_{j_2,M}^{M-1}(z_2) | j_1 \rangle \\ \langle j_4 | \Phi_{\frac{1}{2},1}^0(z_3) \Phi_{j_2,M-1}^M(z_2) | j_1 \rangle & \langle j_4 | \Phi_{\frac{1}{2},1}^1(z_3) \Phi_{j_2,M-1}^{M-1}(z_2) | j_1 \rangle \end{pmatrix} \\ &= x^{-j_2/2\kappa} z_2^{h_4-h_1} g_+(x) \begin{pmatrix} \tilde{\Psi}_{(+)}^0(x) & \tilde{\Psi}_{(+)}^1(x) \\ \tilde{\Psi}_{(+)}^0(x) & \tilde{\Psi}_{(+)}^1(x) \end{pmatrix}, \\ & \begin{pmatrix} \langle j_4 | \Phi_{j_2,M}^M(z_2) \Phi_{\frac{1}{2},0}^0(z_3) | j_1 \rangle & \langle j_4 | \Phi_{j_2,M}^{M-1}(z_2) \Phi_{\frac{1}{2},0}^1(z_3) | j_1 \rangle \\ \langle j_4 | \Phi_{j_2,M-1}^M(z_2) \Phi_{\frac{1}{2},1}^0(z_3) | j_1 \rangle & \langle j_4 | \Phi_{j_2,M-1}^{M-1}(z_2) \Phi_{\frac{1}{2},1}^1(z_3) | j_1 \rangle \end{pmatrix} \\ &= x^{j_2/2\kappa} z_2^{h_4-h_1} g_-(px) \begin{pmatrix} p^{-m+m_1} \tilde{\Psi}_{(-)}^0(px) & p^{-m+m_2} \tilde{\Psi}_{(-)}^0(px) \\ p^{m+m_1} \tilde{\Psi}_{(-)}^0(px) & p^{m+m_2} \tilde{\Psi}_{(-)}^1(px) \end{pmatrix}. \end{aligned} \quad (5.1)$$

Proof. Both sides of the above equations satisfy the same q -KZ equations. So we need only to compare the leading terms. From $g_+(x) = g(x) + O(x^2)$ and $g_-(px) = g(x^{-1}) + O(x^{-2})$, we obtain the required relation. Q.E.D.

From the difference equation for $g(x)$ and $\tilde{\Phi}(x)$ and from the Proposition-I,-II, we have

Theorem. *The exchange relation for the q -vertex operators of spin $1/2$ and arbitrary spin $j_2 \in \frac{1}{2}\mathbf{Z}_{\geq 0}$ is as follows*

$$\check{R}_{ij}^{kl}\left(\frac{z}{w}\right)\Phi_{\frac{1}{2},k}^{\alpha}(w)\Phi_{j_2,l}^{\beta}(z) = \Phi_{j_2,i}^{\gamma}(z)\Phi_{\frac{1}{2},j}^{\delta}(w)\check{C}_{\gamma\delta}^{\alpha\beta}\left(\frac{z}{w}, \lambda\right), \quad (5.2)$$

with

$$\begin{pmatrix} \check{R}_{M,0}^0{}^M(x) & \check{R}_{M,0}^1{}^{M-1}(x) \\ \check{R}_{M-1,1}^0{}^M(x) & \check{R}_{M-1,1}^1{}^{M-1}(x) \end{pmatrix} = \frac{f(x)}{x - q^{M+\overline{M}}} \begin{pmatrix} (xq^M - q^{\overline{M}}) & q^M(q^{-\overline{M}} - q^{\overline{M}}) \\ xq^{\overline{M}}(q^{-M} - q^M) & xq^{\overline{M}} - q^M \end{pmatrix}, \quad (5.3)$$

$$\begin{pmatrix} \check{C}_{M,0}^0{}^M(x, \lambda) & \check{C}_{M,0}^1{}^{M-1}(x, \lambda) \\ \check{C}_{M-1,1}^0{}^M(x, \lambda) & \check{C}_{M-1,1}^1{}^{M-1}(x, \lambda) \end{pmatrix} = x^{-j_2/\kappa} C_g(x) \begin{pmatrix} p^{-m_1} C_0^0(x) & p^{-m_1} C_0^1(x) \\ p^{-m_2} C_1^0(x) & p^{-m_2} C_1^1(x) \end{pmatrix}, \quad (5.4)$$

where $M = \alpha + \beta = \gamma + \delta = i + j = k + l$, $\lambda = 2j_4 + 1$ and $f(x)$, $C_{\beta}^{\alpha}(x)$ and $C_g(x)$ are as given in (4.6), (4.11), and (4.13) respectively.

Proof. From the intertwining property of the R -matrix, it is obvious that the Theorem holds not only for the lowest state $\langle j_4 | \in V_{j_4}^*$ and the highest state $|j_1\rangle \in V_{j_1}$ but also for arbitrary states in $V_{j_4}^*$ and V_{j_1} . Q.E.D.

§ 5.2. If we denote the q -vertex operator and the connection matrix as

$$\begin{aligned} \Phi \begin{bmatrix} j_2 \\ j_3 \ j_1 \end{bmatrix}_{m_2} (z) &= \Phi_{j_2, m_2}^{j_1+j_2-j_3}(z) \quad : \quad V_{j_1} \rightarrow V_{j_3}, \\ C_{\frac{1}{2}j_2} \begin{bmatrix} j_4 \ j' \\ j \ j_1 \end{bmatrix} \left(\frac{z}{w}\right) &= \check{C}_{j'+j_2-j_4, j_1+\frac{1}{2}-j'}^{j+\frac{1}{2}-j_4, j_1+j_2-j} \left(\frac{z}{w}, \lambda\right), \end{aligned} \quad (5.5)$$

then the exchange relation (5.2) can be written as

$$\check{R}_{ij}^{kl}\left(\frac{z}{w}\right)\Phi \begin{bmatrix} \frac{1}{2} \\ j_4 \ j \end{bmatrix}_k (w)\Phi \begin{bmatrix} j_2 \\ j \ j_1 \end{bmatrix}_l (z) = \Phi \begin{bmatrix} j_2 \\ j_4 \ j' \end{bmatrix}_i (z)\Phi \begin{bmatrix} \frac{1}{2} \\ j' \ j_1 \end{bmatrix}_j (w)C_{\frac{1}{2}j_2} \begin{bmatrix} j_4 \ j' \\ j \ j_1 \end{bmatrix} \left(\frac{z}{w}\right). \quad (5.6)$$

From the fact that the R -matrix $\check{R}_{ij}^{kl}(\frac{z}{w})$ satisfies the Yang-Baxter equation, the connection matrix $C_{\frac{1}{2}j_2} \begin{bmatrix} j_4 & j' \\ j & j_1 \end{bmatrix}(\frac{z}{w})$ obeys the IRF type Yang-Baxter relation [4,11]. Moreover if we denote $[n]_{\vartheta} = p^{-n/2\kappa} \Theta(p^{n/\kappa})$, $x = q^{2j_2+1-2u}$, then

$$\begin{pmatrix} \check{C}_{M,0}^{0,M}(x, \lambda) & \check{C}_{M,0}^{1,M-1}(x, \lambda) \\ \check{C}_{M-1,1}^{0,M}(x, \lambda) & \check{C}_{M-1,1}^{1,M-1}(x, \lambda) \end{pmatrix} = \frac{C_g(x)x^{-j_2/\kappa}}{[\lambda + M - \overline{M}]_{\vartheta}[u - M - \overline{M}]_{\vartheta}} \\ \times \begin{pmatrix} \tilde{\alpha}_1 & \\ & \tilde{\alpha}_2 \end{pmatrix} \begin{pmatrix} [\lambda - \overline{M}]_{\vartheta}[u - \overline{M}]_{\vartheta} & [-M]_{\vartheta}[u - \overline{M} + \lambda]_{\vartheta} \\ [\overline{M}]_{\vartheta}[u - M - \lambda]_{\vartheta} & [\lambda + M]_{\vartheta}[u - M]_{\vartheta} \end{pmatrix} \begin{pmatrix} \alpha_1 & \\ & \alpha_2 \end{pmatrix}^{-1}, \quad (5.7)$$

where

$$C_g(x) = \prod_{n \geq 0} \frac{[u - M - \overline{M} - 2n]_{\vartheta}[u - 2 - 2n]_{\vartheta}}{[u - M - \overline{M} - 1 - 2n]_{\vartheta}[u - 1 - 2n]_{\vartheta}}, \quad (5.8)$$

$$\begin{aligned} \alpha_1 &= \frac{\Gamma_p(2m_1)\Gamma_p(2m - m_1 - m_2 + 1)}{\Gamma_p(2m + m_1 - m_2 + 1)} x^{-m-m_1}, \\ \alpha_2 &= \frac{\Gamma_p(2m_2)\Gamma_p(-2m - m_1 - m_2)}{\Gamma_p(-2m - m_1 + m_2)} x^{m-m_2} p^{-m_2}, \\ \tilde{\alpha}_1 &= \frac{\Gamma_p(2m_1)\Gamma_p(-2m - m_1 - m_2)}{\Gamma_p(-2m + m_1 - m_2)} x^{-m+m_1} p^{-m_1}, \\ \tilde{\alpha}_2 &= \frac{\Gamma_p(2m_2)\Gamma_p(2m - m_1 - m_2 + 1)}{\Gamma_p(2m - m_1 + m_2 + 1)} x^{m+m_2}, \end{aligned} \quad (5.9)$$

this connection matrix is equivalent to the elliptic Boltzman weight of IRF type obtained by the fusion procedure in Ref. [13].

6. Conclusion

We have given explicitly the two point functions and the exchange relations for the q -vertex operators whose spins are $1/2$ and arbitrary $j \in \frac{1}{2}\mathbf{Z}_{\geq 0}$. We also confirmed that the connection matrix is equivalent to the elliptic Boltzman weight of IRF type obtained by the fusion procedure in Ref. [13].

We expect that the exchange relation for the two arbitrary spin q -vertex operators will be given by an analogous fusion procedure. Our method essentially relies on the connection formula of q -hypergeometric function and it is applicable only to the case when the number

of intermediate channels is at most two. A more promising approach to the connection problem is to use the integral formula.

Recently, a free field realization for $U_q(\widehat{sl_2})$ was constructed [17,18,19]. The free field realization will give a powerful tool [20] to calculate the integral formulas [1,2,5,6] for the q -KZ solution and to solve their connection problem.

Acknowledgment.

This work has been carried out in collaboration with Y. Yamada. I am grateful to him for help in many ways. I would like to thank M. Jimbo, J. Shiraishi and the members of KEK theory group for valuable discussions. I would also like to thank to N. A. McDougall for a careful reading of the manuscript.

Appendix A. The existence condition for the q -vertex operator of arbitrary spin

§ A.1. Here we briefly review the result of our previous paper [15]. For an arbitrary spin $j \in \mathbf{C}$, the q -vertex operator $\Phi_{j_2}(z) : V_{j_1} \rightarrow V_{j_3}$ is defined by the transformation property under the adjoint action of $U_q(\widehat{sl_2})$ [4,3,14], such that $\text{ad}(a) \Phi(z) \equiv \sum_k a_k^1 \Phi(z) S(a_k^2)$ with $\Delta(a) = \sum_k a_k^1 \otimes a_k^2$.

Definition. The q -vertex operator $\Phi_{j_2}(z) : V_{j_1} \rightarrow V_{j_3}$ is defined as

$$\text{ad}(a) \Phi(z) = \rho(a) \Phi(z), \quad (\text{A.1})$$

for $\forall a \in U_q(\widehat{sl_2})$, where ρ is a certain representation of $U_q(\widehat{sl_2})$.

The following intertwining property is equivalent to above definition;

$$a \Phi(z) = \sum_k \rho(a_k^1) \Phi(z) a_k^2. \quad (\text{A.2})$$

For ρ , if we take the contravariant difference representation defined by

$$\begin{aligned} \rho(e_1) &= x[2j - x \frac{d}{dx}], & \rho(e_0) &= \frac{z}{x} [x \frac{d}{dx}], \\ \rho(k_1) &= q^{2(j-x \frac{d}{dx})}, & \rho(k_0) &= q^{-2(j-x \frac{d}{dx})}, & \rho(d) &= -z \frac{d}{dz}, \\ \rho(f_1) &= \frac{1}{x} [x \frac{d}{dx}], & \rho(f_0) &= \frac{x}{z} [2j - x \frac{d}{dx}], \end{aligned} \quad (\text{A.3})$$

then the q -vertex operator $\Phi_j(z, x)$ of spin j can be defined explicitly as follows [15]

$$\begin{aligned} e_i \Phi_j(z, x) &= \rho(e_i) \Phi_j(z, x) k_i + \Phi_j(z, x) e_i, \\ k_i \Phi_j(z, x) &= \rho(k_i) \Phi_j(z, x) k_i, \\ f_i \Phi_j(z, x) &= \rho(f_i) \Phi_j(z, x) + \rho(k_i^{-1}) \Phi_j(z, x) f_i, \end{aligned} \tag{A.4}$$

and

$$d \Phi_j(z, x) = \rho(d) \Phi_j(z, x) + \Phi_j(z, x) d. \tag{A.5}$$

Here $[n]$ denotes the q integer $[n] = (q^n - q^{-n})/(q - q^{-1})$, so the $\rho(J)$'s are the difference operators, e.g. $q^x \frac{d}{dx} f(x) = f(qx)$ for any function $f(x)$.

And we normalize the ground state matrix element of the q -vertex operator as

$$\langle j_3 | \Phi_{j_2}(z, x) | j_1 \rangle = z^{h_3 - h_1} x^{j_1 + j_2 - j_3}. \tag{A.6}$$

For $2j+1 \in \mathbf{Z}_{>0}$, if we set $\Phi_j(z, x) = \sum_{m=0}^{2j} \Phi_{j,m}(z) x^m$, then we have the previous definition for the q -vertex operators $\Phi_{j,m}(z)$ in §3.1.

§ **A.2.** The q -vertex operator $\Phi_{j_2}(z, x) : V_{j_1} \rightarrow V_{j_3}$ exists if and only if

$$\langle j_3 | \Phi_{j_2}(z, x) | \chi_1 \rangle = \langle \chi_3 | \Phi_{j_2}(z, x) | j_1 \rangle = 0, \tag{A.7}$$

for all the null vectors $|\chi_1\rangle \in V_{j_1}$ and $\langle \chi_3 | \in V_{j_3}^*$.

For $r, s \in \mathbf{Z}$ and $j_n \in \mathbf{C}$, let

$$\begin{aligned} f_{r,s}(j_1, j_2, j_3) &= \prod_{n=0}^{r-1} \prod_{m=0}^s [j_1 + j_2 - j_3 - n + m\kappa] \prod_{n=1}^r \prod_{m=1}^s [-j_1 + j_2 + j_3 + n - m\kappa], \\ f_{r,s}(j_1, j_2, j_3) &= \prod_{n=0}^{-r-1} \prod_{m=0}^{-s-1} [-j_1 + j_2 + j_3 - n + m\kappa] \prod_{n=1}^{-r} \prod_{m=1}^{-s-1} [j_1 + j_2 - j_3 + n - m\kappa]. \end{aligned} \tag{A.8}$$

for the case (i) $r > 0$ and $s \geq 0$ or (ii) $r < 0$ and $s < 0$, respectively. Then we have

Theorem.[15] *The existence conditions for the q -vertex operators $\Phi_{j_2}(z, x) : V_{j_1} \rightarrow V_{j_3}$ are given as follows.*

(I). For the rational level $\kappa = p/q$, with the coprime integers p and q , and $2j_n + 1 = r_n - s_n \kappa$ with $0 < r_n < p$ and $0 \leq s_n < q$, ($n = 1, 2, 3$), the q -vertex operator exists if and only if

$$f_{r_1, s_1}(j_1, j_2, j_3) = f_{r_1 - p, s_1 - q}(j_1, j_2, j_3) = 0, \quad (\text{A.9})$$

(II). For the generic level, the q -vertex operator exists if and only if

$$\sum_{r_1, s_1 \in \mathbf{Z}} f_{r_1, s_1}(j_1, j_2, j_3) \delta_{r_1 - s_1 \kappa}^{2j_1 + 1} = \sum_{r_3, s_3 \in \mathbf{Z}} f_{r_3, s_3}(j_3, j_2, j_1) \delta_{r_3 - s_3 \kappa}^{2j_3 + 1} = 0. \quad (\text{A.10})$$

Proof is given by using the following Lemma-I and Lemma-II.

Lemma I.[21] For $\kappa = k + 2 \in \mathbf{C} \setminus \{0\}$ and the highest weight j , parametrized as $2j_{r,s} + 1 = r - s\kappa$ with $r, s \in \mathbf{Z}$, such that (i) $r > 0$ and $s \geq 0$ or (ii) $r < 0$ and $s < 0$, there exists a unique null vector $|\chi_{r,s}\rangle \in V_j$ of grade $N = rs$ and charge $Q = -r$. And the null vector in $V_{j_{r,s}}$ is as follows,

$$\begin{aligned} |\chi_{r,s}\rangle &= (f_1)^{r+s\kappa} (f_0)^{r+(s-1)\kappa} \dots \dots (f_0)^{r-(s-1)\kappa} (f_1)^{r-s\kappa} |j_{r,s}\rangle, \\ |\chi_{r,s}\rangle &= (f_0)^{-r-(s+1)\kappa} (f_1)^{-r-(s+2)\kappa} \dots \dots (f_1)^{-r+(s+2)\kappa} (f_0)^{-r+(s+1)\kappa} |j_{r,s}\rangle, \end{aligned} \quad (\text{A.11})$$

for the cases (i) and (ii) respectively.

Lemma II. For the null vectors $|\chi_{r,s}\rangle \in V_{j_1}$ and $\langle \chi_{r,s} | \in V_{j_3}^*$, we have

$$\begin{aligned} \langle j_3 | \Phi_{j_2}(z, x) | \chi_{r,s} \rangle &= f_{r,s}(j_1, j_2, j_3) z^{h_3 - h_1 - rs} x^{j_1 + j_2 - j_3 - r}, \\ \langle \chi_{r,s} | \Phi_{j_2}(z, x) | j_1 \rangle &= f_{r,s}(j_3, j_2, j_1) z^{h_3 - h_1 + rs} x^{j_1 + j_2 - j_3 + r}, \end{aligned} \quad (\text{A.12})$$

up to some non-zero multiple factors.

Appendix B. Calculation of the two point function by the q -OPE

§ B.1. We calculate the image $\Phi_{j_2, m_2}(z) |j_1\rangle \in V_{j_3}$ of the highest weight vector $|j_1\rangle$. Let

$$\Phi_{j_2, m_2}(z) |j_1\rangle = \delta_{m_2}^{j_3 - j_1 + Q} \sum_N z^{h_3 - h_1 + N} q^{kN + 2j_1 Q} |N, Q\rangle_{j_3}, \quad (\text{B.1})$$

where $|N, Q\rangle_{j_3}$ is the homogeneous components of grade N and charge $-Q$, such that

$$|N, Q\rangle_{j_3} = \sum_{(\alpha_1, \dots, \alpha_n)} \beta_{\alpha_1, \dots, \alpha_n} f_{\alpha_1 \dots \alpha_n} |j_3\rangle, \quad (\text{B.2})$$

$N = \sum_i \bar{\alpha}_i$, $Q = \sum_i (\alpha_i - \bar{\alpha}_i)$, $\alpha_i = 0, 1$, $\bar{\alpha} = 1 - \alpha$ and $f_{\alpha_1 \dots \alpha_n} = f_{\alpha_1} \dots f_{\alpha_n}$. From the definition of the q -vertex operator (3.1), we have the descent equations for $|N, Q\rangle_{j_3}$ [15]

$$\begin{aligned} e_1 |N, Q\rangle_{j_3} &= [-j_1 + j_2 + j_3 - Q + 1] |N, Q - 1\rangle_{j_3}, \\ e_0 |N, Q\rangle_{j_3} &= [j_1 + j_2 - j_3 + Q + 1] |N - 1, Q + 1\rangle_{j_3}. \end{aligned} \quad (\text{B.3})$$

From these descent equations, we can calculate the expansion coefficients $\beta_{\alpha_1, \dots, \alpha_n}$.

Example. The q -OPE of spin $1/2$ q -vertex operator is

$$\begin{aligned} \Phi_{\frac{1}{2}, 0}^0(z) |j\rangle &= z^{h^+ - h} (1 + q^k z (\beta_{01}^+ f_{01} + \beta_{10}^+ f_{10}) + O(z^2)) |j + \frac{1}{2}\rangle, \\ \Phi_{\frac{1}{2}, 1}^0(z) |j\rangle &= z^{h^+ - h} q^{2j} (\beta_1^+ f_1 + q^k z (\beta_{011}^+ f_{011} + \beta_{101}^+ f_{101} + \beta_{110}^+ f_{110}) + O(z^2)) |j + \frac{1}{2}\rangle, \\ \Phi_{\frac{1}{2}, 1}^1(z) |j\rangle &= z^{h^- - h} (1 + q^k z (\beta_{10}^- f_{10} + \beta_{01}^- f_{01}) + O(z^2)) |j - \frac{1}{2}\rangle, \\ \Phi_{\frac{1}{2}, 0}^1(z) |j\rangle &= z^{1+h^- - h} q^{k-2j} (\beta_0^- f_0 + q^k z (\beta_{100}^- f_{100} + \beta_{010}^- f_{010} + \beta_{001}^- f_{001}) + O(z^2)) |j - \frac{1}{2}\rangle, \end{aligned} \quad (\text{B.4})$$

where $h = h_j$, $h^\pm = h_{j \pm \frac{1}{2}}$,

$$\begin{aligned} \beta_1^+ &= \frac{1}{[2j+1]}, & \beta_{011}^+ &= \frac{-[k+2][2j+2]}{[k+2j+3][2][k+2][2j+1]}, \\ \beta_{01}^+ &= \frac{[2j+3]}{[2][k+2][2j+1]}, & \beta_{101}^+ &= \frac{[k+2][2j] + [k+3][2j+3]}{[k+2j+3][2][k+2][2j+1]}, \\ \beta_{10}^+ &= \frac{-1}{[2][k+2]}, & \beta_{110}^+ &= \frac{-[k+3]}{[k+2j+3][2][k+2]}, \end{aligned} \quad (\text{B.5})$$

and

$$\beta_{\alpha_1, \dots, \alpha_n}^-(2j) = \beta_{\bar{\alpha}_1, \dots, \bar{\alpha}_n}^+(k-2j). \quad (\text{B.6})$$

§ B.2. By using the q -OPE and one point function, we can in principle derive the arbitrary N point functions.

Example. The two point functions for the spin $\frac{1}{2}$ and arbitrary spin j_2 q -vertex operators are

$$\begin{aligned}
\langle j_4 | \Phi_{\frac{1}{2},0}^0(z_3) \Phi_{j_2,M}^M(z_2) | j_1 \rangle &= z_3^{h_4-h_4^-} z_2^{h_4^- - h_1} \\
&\times \left\{ 1 + \frac{q^{k+2}x}{[2][k+2][k-2j_4+1]} \begin{pmatrix} [k-2j_4+3][M][2j_2-M+1] \\ -[k-2j_4+1][M+1][2j_2-M] \end{pmatrix} + O(x^2) \right\}, \\
\langle j_4 | \Phi_{\frac{1}{2},1}^0(z_3) \Phi_{j_2,M-1}^M(z_2) | j_1 \rangle &= -z_3^{h_4-h_4^-} z_2^{h_4^- - h_1} x q^{k-2j_1+1} \frac{[M]}{[k-2j_4+1]} \\
&\times \left\{ 1 + \frac{q^{k+2}x}{[2][k+2][2k-2j_4+3]} \begin{pmatrix} -[k+2][k-2j_4+2][M-1][2j_2-M+2] \\ +[k+2][k-2j_4][M][2j_2-M+1] \\ +[k+3][k-2j_4+3][M][2j_2-M+1] \\ -[k+3][k-2j_4+1][M+1][2j_2-M] \end{pmatrix} + O(x^2) \right\},
\end{aligned} \tag{B.7}$$

$$\begin{aligned}
\langle j_4 | \Phi_{\frac{1}{2},0}^1(z_3) \Phi_{j_2,M}^{M-1}(z_2) | j_1 \rangle &= -z_3^{h_4-h_4^+} z_2^{h_4^+ - h_1} q^{2j_1+1} \frac{[2j_2-M+1]}{[2j_4+1]} \\
&\times \left\{ 1 + \frac{q^{k+2}x}{[2][k+2][k+2j_4+3]} \begin{pmatrix} -[k+2][2j_4+2][M+1][2j_2-M] \\ +[k+2][2j_4][M][2j_2-M+1] \\ +[k+3][2j_4+3][M][2j_2-M+1] \\ -[k+3][2j_4+1][M-1][2j_2-M+2] \end{pmatrix} + O(x^2) \right\}, \\
\langle j_4 | \Phi_{\frac{1}{2},1}^1(z_3) \Phi_{j_2,M-1}^{M-1}(z_2) | j_1 \rangle &= z_3^{h_4-h_4^+} z_2^{h_4^+ - h_1} \\
&\times \left\{ 1 + \frac{q^{k+2}x}{[2][k+2][2j_4+1]} \begin{pmatrix} [2j_4+3][M][2j_2-M+1] \\ -[2j_4+1][M-1][2j_2-M+2] \end{pmatrix} + O(x^2) \right\},
\end{aligned} \tag{B.8}$$

and

$$\begin{aligned}
\langle j_4 | \Phi_{j_2,M}^M(z_2) \Phi_{\frac{1}{2},0}^0(z_3) | j_1 \rangle &= z_2^{h_4-h_1^+} z_3^{h_1^+ - h_1} \\
&\times \left\{ 1 + \frac{q^{k+2}x^{-1}}{[2][k+2][2j_1+1]} \begin{pmatrix} [2j_1+3][2j_2-M+1][M] \\ -[2j_1+1][2j_2-M][M+1] \end{pmatrix} + O\left(\frac{1}{x^2}\right) \right\}, \\
\langle j_4 | \Phi_{j_2,M-1}^M(z_2) \Phi_{\frac{1}{2},1}^0(z_3) | j_1 \rangle &= -z_2^{h_4-h_1^+} z_3^{h_1^+ - h_1} q^{2j_4+1} \frac{[M]}{[2j_1+1]} \\
&\times \left\{ 1 + \frac{q^{k+2}x^{-1}}{[2][k+2][k+2j_1+3]} \begin{pmatrix} -[k+2][2j_1+2][2j_2-M+2][M-1] \\ +[k+2][2j_1][2j_2-M+1][M] \\ +[k+3][2j_1+3][2j_2-M+1][M] \\ -[k+3][2j_1+1][2j_2-M][M+1] \end{pmatrix} + O\left(\frac{1}{x^2}\right) \right\},
\end{aligned} \tag{B.9}$$

$$\begin{aligned}
\langle j_4 | \Phi_{j_2, M}^{M-1}(z_2) \Phi_{\frac{1}{2}, 0}^1(z_3) | j_1 \rangle &= -z_2^{h_4 - h_1^-} z_3^{h_1^- - h_1} x^{-1} q^{k-2j_4+1} \frac{[2j_2 - M + 1]}{[k - 2j_1 + 1]} \\
&\times \left\{ 1 + \frac{q^{k+2} x^{-1}}{[2][k+2][2k-2j_1+3]} \begin{pmatrix} -[k+2][k-2j_1+2][2j_2-M][M+1] \\ +[k+2][k-2j_1][2j_2-M+1][M] \\ +[k+3][k-2j_1+3][2j_2-M+1][M] \\ -[k+3][k-2j_1+1][2j_2-M+2][M-1] \end{pmatrix} \right\} + O\left(\frac{1}{x^2}\right), \\
\langle j_4 | \Phi_{j_2, M-1}^{M-1}(z_2) \Phi_{\frac{1}{2}, 1}^1(z_3) | j_1 \rangle &= z_2^{h_4 - h_1^-} z_3^{h_1^- - h_1} \\
&\times \left\{ 1 + \frac{q^{k+2} x^{-1}}{[2][k+2][k-2j_1+1]} \begin{pmatrix} [k-2j_1+3][2j_2-M+1][M] \\ -[k-2j_1+1][2j_2-M+2][M-1] \end{pmatrix} \right\} + O\left(\frac{1}{x^2}\right),
\end{aligned} \tag{B.10}$$

where $x = z_2/z_3$, $h_n^\pm = h_{j_n \pm \frac{1}{2}}$ and $M = j_1 + j_2 - j_4 + \frac{1}{2}$.

Appendix C. The R -matrix

For the finite dimensional representation $V_j(z)$ in §2.2, let $|M_1, M_2\rangle = |1/2, M_1\rangle \otimes |j, M_2\rangle \in V_{1/2}(z) \otimes V_j(w)$. The R -matrix, $R(z/w) : V_{1/2}(z) \otimes V_j(w) \rightarrow V_{1/2}(z) \otimes V_j(w)$, defined by $R(z/w)\Delta'_{z,w} = \Delta_{z,w}R(z/w)$, is a 2×2 block diagonal form in each sector $\mathbf{C}|0, M\rangle \oplus \mathbf{C}|1, M-1\rangle$, with $M \in \{1, \dots, 2j-1\}$.

The R -matrix $\tilde{R}(x)$, which is normalized as $\tilde{R}(x)|0, 0\rangle = |0, 0\rangle$, is given explicitly as follows

$$\begin{pmatrix} \tilde{R}(x)|0, M\rangle \\ \tilde{R}(x)|1, M-1\rangle \end{pmatrix} = \begin{pmatrix} a_M(x) & b_M(x) \\ c_M(x) & d_M(x) \end{pmatrix} \begin{pmatrix} |0, M\rangle \\ |1, M-1\rangle \end{pmatrix}, \tag{C.1}$$

where

$$\begin{pmatrix} a_M(x) & b_M(x) \\ c_M(x) & d_M(x) \end{pmatrix} = \frac{1}{x - q^{M+\overline{M}}} \begin{pmatrix} xq^M - q^{\overline{M}} & q^M(q^{-\overline{M}} - q^{\overline{M}}) \\ xq^{\overline{M}}(q^{-M} - q^M) & xq^{\overline{M}} - q^M \end{pmatrix}, \tag{C.2}$$

with $\overline{M} = 2j + 1 - M$. They obey the following crossing relation

$$\begin{aligned}
(((\tilde{R}(x)^{-1})^{t_1})^{-1})^{t_1} &= \frac{(1 - xq^{-2j+3})(1 - xq^{2j+1})}{(1 - xq^{-2j+1})(1 - xq^{2j+3})} K^{-1} \tilde{R}(xq^{-4}) K, \\
K &= \begin{pmatrix} q & \\ & q^{-1} \end{pmatrix} \otimes 1,
\end{aligned} \tag{C.3}$$

where t_1 means a transpose on the first component.

The R -matrix $R(x)$ which is the image of the universal R -matrix can be determined by the crossing relation [4]

$$(((R(x)^{-1})^{t_1})^{-1})^{t_1} = K^{-1}R(xq^{-4})K. \quad (\text{C.4})$$

The relation between R and $\tilde{R}$ is then given as

$$R(x) = f(x)\tilde{R}(x), \quad f(x) = \frac{(1 - xq^{2j_2+3})(1 - xq^{-2j_2+1})}{(1 - xq^{2j_2+1})(1 - xq^{-2j_2+3})}f(xq^4). \quad (\text{C.5})$$

The solution for $f(x)$ in the region $x \ll 1$ is

$$f(x) = \prod_{n=0}^{\infty} \frac{(1 - xq^{2j+3+4n})(1 - xq^{-2j+1+4n})}{(1 - xq^{2j+1+4n})(1 - xq^{-2j+3+4n})}. \quad (\text{C.6})$$

This solution can be analytically continued to the whole $x \in \mathbf{C}$ uniquely.

References

- [1] K. Aomoto, “ A note on holonomic q -difference systems ” Algebraic Analysis I, ed. by M. Kashiwara and T. Kawai, Academic Press, San Diego (1988) 25-28.
- [2] K. Mimachi, “ Connection problem in holonomic q -difference system associated with a Jackson integral of Jordan-Pochhammer type ” Nagoya Math. Journ., **116** (1989) 149-161.
- [3] F.A. Smirnov, “ Dynamical symmetries of massive integrable models ” Int. J. Mod. Phys. **A7** Supplement 1 (1992) 813-837;839-818.
- [4] I.B. Frenkel and N.Yu. Reshetikhin, “ Quantum affine algebras and holonomic difference equations ” Commun. Math. Phys. **146** (1992) 1-60.
- [5] A. Matsuo, “ Jackson integrals of Jordan-Pochhammer typ and quantum Knizhnik-Zamolodchikov equations ” preprint (1991); “ Quantum algebra structure of certain Jackson integrals ” preprint (1992).
- [6] N. Reshetikhin, “ Jackson-type integrals, Bethe vectors, and solutions to a difference analog of the Knizhnik-Zamokodchikov system ” preprint (1991).
- [7] K. Aomoto, Y. Kato and K. Mimachi, “ A solution of Yang-Baxter equation as connection coefficidnts of a holonomic q -difference system ” Duke Math. Jour. **65** IMRN (1992) 7-15.
- [8] B. Davies, O. Foda, M. Jimbo, T. Miwa and A. Nakayashiki, “ Diagonalization of the XXZ hamiltonian by vertex operators ” RIMS preprint (1992).
- [9] M. Idzumi, K. Iohara, M. Jimbo, T. Miwa, T. Nakashima and T. Tokihiro, “ Quantum affine symmetry in vertex models ” RIMS preprint (1992).
- [10] A. Tsuchiya and Y. Kanie, “ Vertex operators in conformal field theory on P^1 and monodromy representations of braid group ” Advanced Studies in Pure Math. **16** (1988) 297-372, Lett. Math. Phys. **13** (1987) 303.
- [11] E. Date, M. Jimbo and M. Okado, “ Crystal base and q -vertex operators ” Osaka Univ. Math. Sci. preprint 1 (1991).
- [12] G.E. Andrews, R.J. Baxter and P.J. Forrester, “ Eight-vertex SOS model and generalized Rogers-Ramanujan-type identities ” Jour. Stat. Phys. **35** (1984) 193-266.
- [13] E. Date, M. Jimbo, T. Miwa and M. Okado, “ Solvable lattice models ” RIMS Preprint 590 (1987).
- [14] G. Felder and A. LeClair, “ Restricted quantum affine symmetry of perturbed minimal conformal models ” Int. J. Mod. Phys. **A7** (1992) 239-278.
- [15] H. Awata and Y. Yamada, “ Fusion rules for the q -vertex operators of $U_q(\widehat{sl_2})$ ” Mod. Phys. Lett. **A7** (1992) 2235-2243.
- [16] H. Awata and Y. Yamada, “ Fusion rules for the fractional level $\widehat{sl(2)}$ algebra ” Mod. Phys. Lett. **A7** (1992) 1185-1195.

- [17] A. Matsuo, “ Free field representation of quantum affine algebra $U_q(\widehat{sl_2})$ ” preprint (1992).
- [18] A. Abada, A.H. Bougourzi and M.A. El Gradechi, “ Deformation of the Wakimoto construction ” preprint (1992).
- [19] J. Shiraishi, “ Free boson representation of $U_q(\widehat{sl_2})$ ” preprint UT-617 (1992).
- [20] H. Awata, A. Tsuchiya and Y. Yamada, “ Integral formulas for the WZNW correlation functions ” Nucl. Phys. **B365** (1991) 680-696.
- [21] F.G. Malikov, “ Quantum groups: Singular vectors and BGG resolution ” RIMS Preprint 835 (1991).